



University of Tennessee, Knoxville
**TRACE: Tennessee Research and Creative
Exchange**

Faculty Publications and Other Works --
Biomedical and Diagnostic Sciences

Veterinary Medicine -- Faculty Publications and
Other Works

2014

Pharmacokinetics of intravenous and subcutaneous ceftiofur in alpacas

S COX

University of Tennessee, Knoxville

C. SOMMARDAHL

University of Tennessee, Knoxville

R. SEDDIGHI

University of Tennessee, Knoxville

R. VIDELA

University of Tennessee, Knoxville

J. HAYES

University of Tennessee, Knoxville

See next page for additional authors

Follow this and additional works at: https://trace.tennessee.edu/utk_compmedpubs

Recommended Citation

COX, S; SOMMARDAHL, C.; SEDDIGHI, R.; VIDELA, R.; HAYES, J.; nPISTOLE, N.; HAMIL, M.; and DOHERTY, T., "Pharmacokinetics of intravenous and subcutaneous ceftiofur in alpacas" (2014). *Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences*.
https://trace.tennessee.edu/utk_compmedpubs/145

This Article is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Authors

S COX, C. SOMMARDAHL, R. SEDDIGHI, R. VIDELA, J. HAYES, N. nPISTOLE, M. HAMIL, and T. DOHERTY

J. vet. Pharmacol. Therap. 38, 344–349. doi: 10.1111/jvp.12181.

Pharmacokinetics of intravenous and subcutaneous ceftiofur in alpacas

S. COX*

C. SOMMARD AHL†

R. SEDDIGHI†

R. VIDELA†

J. HAYES*

N. PISTOLE*

M. HAMILL* &

T. DOHERTY†

Cox, S., Sommardahl, C., Seddighi, R., Videla, R., Hayes, J., Pistole, N., Hamill, M., Doherty, T. Pharmacokinetics of intravenous and subcutaneous ceftiofur in alpacas. *J. vet. Pharmacol. Therap.* 38, 344–349.

The purpose of this study was to determine the pharmacokinetics of ceftiofur after intravenous and subcutaneous dose of 8 mg/kg to alpacas. Bacterial infections requiring long-term antibiotic therapy such as neonatal bacteraemia, pneumonia, peritonitis, dental, and uterine infections are a significant cause of morbidity and mortality in this species. However, few antimicrobials have been evaluated and proven to have favorable pharmacokinetics for therapeutic use. Most antimicrobials that are currently used require daily injections for many days. Ceftiofur is a long-acting cephalosporin that is formulated for subcutaneous administration, and its long-elimination half-life allows for 14-day dosing intervals in dogs and cats. The properties of ceftiofur may be advantageous for medical treatment of camelids due to its broad spectrum, route of administration, and long duration of activity. Pharmacokinetic evaluation of antimicrobial drugs in camelids is essential for the proper treatment and prevention of bacterial disease, and to minimize development of antibiotic resistant bacterial strains due to inadequate antibiotic concentra-

*Department of Biomedical and Diagnostic Sciences, College of Veterinary Medicine, The University of Tennessee, Knoxville, TN, USA; †Large Animal Clinical Sciences, College of Veterinary Medicine, The University of Tennessee, Knoxville, TN, USA